



STRUCTURAL EVOLUTION OF THE ANTI-ATLAS DOMAIN :AN OVERVIEW pour Yves Missenard

STRUCTURAL EVOLUTION OF THE ANTI-ATLAS DOMAIN :AN OVERVIEW pour Yves Missenard The Anti-Atlas represents the most important segments of the major Pan-African ($\approx 0.5\text{Ga}$) belt system of North Africa. This orogen exposed in a series of sporadic SW–NE trending outcrops over 700km across southern Morocco, and reach $\approx 150\text{km}$ wide in the central part, west of Ouarzazate (Figure G1 & G2). Scattered outcrops of related rocks are described in NW Algeria which indicate that the belt continues from SE Morocco southeastwards with a NW–SE trend. The Anti-Atlas orogen comprises two main sequences of rocks: a metamorphic basement of Palaeoproterozoic (2Ga) age and the Neoproterozoic rocks. The Palaeoproterozoic rocks form the northern margin of the West African Craton mainly outcropping in the Reguibate Shield in Mauritania And Algeria. The Palaeoproterozoic basement is exposed in a series of uplifted inliers ("boutonniers") surrounded by the Neoproterozoic rocks that were locally deformed with the basement during the Pan African Orogeny. with the during the Pan African Orogeny. Paleoproterozoic rocks and Eburnian orogeny The oldest rocks of Morocco, Archean in age 3 Ga (Montero et al, 2014) crop out in the southernmost Reguibat E shield which form the northern part the West African Craton (WAC). Further north, in the Anti-Atlas chain, the basement units are Paleoproterozoic (Taznakht Group, former "PI") cropping

only in the south of the Anti-Atlas Major Fault (AAMF). They are metamorphic rocks (greenschist to amphibolite facies) intruded by peraluminous and calc-alkaline granitoids dated around 2Ga (Thomas et al, 2002). The corresponding tectono-magmatic and metamorphic events are assigned to the Eburnian/Birimian orogeny. Neoproterozoic rocks and Pan-African orogeny As everywhere in the WAC, Mesoproterozoic rocks are lacking in the Anti-Atlas. Neoproterozoic Formations overlie directly the Paleoproterozoic basement. The Neoproterozoic/Paleoproterozoic boundary is generally a tectonic contact (thrust, strike-slip or detachment faults), and seldom a stratigraphic contact (Tizi n'Taghatine). Globally, the Neoproterozoic formations record three main stages of the Pan-African cycle (figure G4): I. The early Neoproterozoic platform development is marked by the accumulation of thousands of meters of quartzites and stromatolitic limestones (Taghdoute Group, former "PII"), intruded by doleritic dykes and gabbroic intrusions. These rocks are associated with the rifting of the WAC margin, broadly contemporaneous with the oceanic accretion further north (760 Ma), witnessed along the AAMF by the Bou Azzer-El Graara and Siroua ophiolitic sequences (Bou Azzer Group). II. Oceanic closure and subsequent Pan-African collision are associated with oceanic subduction along the northern margin of the WAC, ending with the accretion of oceanic arc formations (figure G4). The reported "blueschist facies" mineral associations in the Bou Azzer inlier are controversial. The polarity of the subduction remains also matter of debates; the same is true for the real location of the northern edge of the WAC. The oblique Pan-African collision (655 Ma to 640 Ma) generated south-verging thrust sheets onto the cratonic margin. South of the AAMF, the main Pan-African phase is recorded by low grade recrystallizations, synmetamorphic folds and various ductile and brittle structures in the Taghdout Group series. III. The Late to Post Pan-African extensional event is recorded by the extensive volcanic and volcano clastic series of the Ouarzazate Group (former "PIII", 580 Ma to 560 Ma), interbedded with subaerial to lacustrine deposits, which unconformably overlie the Eburnian and/or Pan-African basement units. The Ouarzazate Group shows abrupt variations of thickness and facies controlled by extensional tectonic activity. Various high-K calc-alkaline to alkaline plutons emplaced within the Ouarzazate Group, coeval with the volcanic rocks of comparable chemistry. Figure G4: Generalized lithostratigraphic column for the Anti-Atlas Pan-African orogen. "PI" = "XI", etc. are the classical stratigraphic symbols used on Anti-Atlas geological maps. HKCA: High-K calc-alkaline (granitoids) after Thomas et al. (2004), Gasquet et al. (2005) and Liégeois et al. 2006 in Youbi et al., 2013
 Source web par societadageologica.es